

INELASTIC ELECTRON PHOTON SCATTERING AT MODERATE FOUR-MOMENTUM TRANSFERS

PLUTO Collaboration

Ch. BERGER, H. GENZEL, R. GRIGULL, W. LACKAS and F. RAUPACH

I. Physikalisches Institut der RWTH Aachen¹, Germany

A. KLOVNING, E. LILLESTÖL and J.A. SKARD

University of Bergen², Norway

H. ACKERMANN, J. BÜRGER, L. CRIEGEE, H.C. DEHNE, A. ESKREYS³, G. FRANKE,
W. GABRIEL⁴, Ch. GERKE, G. KNIES, E. LEHMANN, H.D. MERTIENS, U. MICHELSEN,
K.H. PAPE, H.D. REICH, M. SCARR⁵, B. STELLA⁶, U. TIMM, W. WAGNER,
P. WALOSCHEK, G.G. WINTER and W. ZIMMERMANN

Deutsches Elektronen Synchrotron DESY, Hamburg, Germany

O. ACHTERBERG, V. BLOBEL⁷, L. BOESTEN, V. HEPP⁸, H. KAPITZA, B. KOPPITZ,
B. LEWENDEL, W. LÜHRSEN, R. van STAA and H. SPITZER

II. Institut für Experimentalphysik der Universität Hamburg¹, Germany

C.Y. CHANG, R.G. GLASSER, R.G. KELLOGG, K.H. LAU, R.O. POLVADO,
B. SECHI-ZORN, A. SKUJA, G. WELCH and G.T. ZORN

University of Maryland⁹, USA

A. BÄCKER¹⁰, F. BARREIRO, S. BRANDT, K. DERIKUM, C. GRUPEN, H.J. MEYER,
B. NEUMANN, M. ROST and G. ZECH

Gesamthochschule Siegen¹, Germany

H.J. DAUM, H. MEYER, O. MEYER, M. RÖSSLER and D. SCHMIDT

Gesamthochschule Wuppertal¹, Germany

Received 9 October 1980

We present new high statistics data on hadron production in photon-photon reactions. The data are analyzed in terms of an electron-photon scattering formalism. The dependence of the total cross section on Q^2 , the four-momentum transfer squared of the scattered electron, and on the mass W of the hadronic system is investigated. The data are compared to predictions from Vector-Meson Dominance and the quark model.

¹ Supported by the BMFT, Germany.

² Partially supported by the Norwegian Council for Science and Humanities.

³ Now at Institute of Nuclear Physics, Krakow, Poland.

⁴ Now at Max Planck Institut für Limnologie, Plön, Germany.

⁵ On leave from University of Glasgow, UK.

⁶ On leave from University of Rome, Italy; partially supported by INFN.

⁷ Now at CERN, Geneva, Switzerland.

⁸ Now at Heidelberg University, Germany.

⁹ Partially supported by Department of Energy, USA.

¹⁰ Now at Harvard University, USA.

High energy electron–positron storage rings offer the opportunity to measure the inclusive reaction $e^+e^- \rightarrow e^+e^- + \text{hadrons}$. These reactions are usually interpreted as hadron production by two virtual photons which are radiated from the incoming leptons ($e^+e^- \rightarrow e^+e^- \gamma^* \gamma^* \rightarrow e^+e^- + \text{hadrons}$). Experimentally these reactions can be easily separated from the annihilation processes ($e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons}$) by detecting one of the outgoing leptons at small angles in the forward direction (“single tagging”). If on the other hand the scattering angle of the second lepton is kept very small by excluding events with an angle greater than e.g. 25 mrad (“antitagging”), the radiated photon is practically on mass shell and purely transverse polarized. This is the kinematical situation of electron scattering (lepton 1 in ref. [1]) off a real photon beam.

The inclusive cross section for this process ($e\gamma \rightarrow eX$) is given by:

$$d\sigma(e\gamma \rightarrow eX) = \Gamma_t [\sigma_t(Q^2, W) + \epsilon \sigma_q(Q^2, W)] d\Omega_1 dE'_1. \quad (1)$$

The kinematical quantities are explained in ref. [1]. Γ_t is the flux factor for the incoming virtual photons radiated from lepton 1 and ϵ is a polarization parameter for the virtual photons. σ_t is the total cross section for the scattering of virtual transverse polarized photons off real photons and σ_q the respective cross section for virtual longitudinal photons. Γ_t and ϵ are respectively given by:

$$\Gamma_t = \alpha E'_1 [1 + (1 - y)^2] / 2\pi^2 Q^2 y, \quad (2)$$

$$\epsilon = 2(1 - y) / [1 + (1 - y)^2],$$

with

$$y = 1 - (E'_1/E) \cos^2(\Theta_1/2) \quad (3)$$

as a measure for the relative energy of the virtual photon. The inclusive cross section for the reaction $e^+e^- \rightarrow e^+e^- + \text{hadrons}$ is then given by:

$$\begin{aligned} \sigma(e^+e^- \rightarrow e^+e^- + \text{hadrons}) \\ = d\sigma(e\gamma \rightarrow eX) N(z, \Theta_{2\max}) dz. \end{aligned} \quad (4)$$

$N(z, \Theta_{2\max}) dz$ is the number of incoming quasireal photons in the energy interval dz ($z = E_\gamma/E$) radiated from leptons with a maximum scattering angle $\Theta_{2\max}$. A standard expression is given by [2,3]:

$$\begin{aligned} N(z, \Theta_{2\max}) = (\alpha/\pi z) \{ [1 + (1 - z)^2] \\ \times \ln \{ (E/m) [(1 - z)/z] \Theta_{2\max} \} - 1 + z \}. \end{aligned} \quad (5)$$

A similar formula published in the literature produces numerical differences of less than 5% for the experimental conditions accessible to our detector. The cross section definition for photon–photon reactions (σ_t, σ_q) agrees with ref. [1] and leads to the above expressions for Γ_t and ϵ which slightly differ from those given in our first paper [5], but do not contain any small angle approximations. $\Gamma_t N$ exactly reproduces the single tag luminosity function given in formula (11) of ref. [6].

In our previous publication [5], we presented cross section measurements up to values of W of 6.5 GeV at $\langle Q^2 \rangle = 0.1 \text{ GeV}^2$. Our new data extend this range up to $W = 8.5 \text{ GeV}$ and $Q^2 = 0.6 \text{ GeV}^2$. The much higher number of events allows for a detailed study of the W and Q^2 dependence.

The experiment has been performed using the detector PLUTO at PETRA, the e^+e^- colliding beam machine at DESY. A description of the experimental set up can be found in refs. [5] and [7]. The data reported in this paper have been taken at beam energies between 15 and 16 GeV ($\langle E \rangle = 15.5 \text{ GeV}$) for an integrated luminosity of 2700 nb^{-1} as determined by the luminosity monitor.

Hadronic events from electron–photon scattering have been defined by the following conditions:

(a) A tag signal on one side, i.e. we have requested a deposited energy of more than 4 GeV in one of the small angle taggers (SAT) of the forward spectrometers ($23 \text{ mrad} < \Theta_1 < 55 \text{ mrad}$).

(b) “Antitagging” on the opposite side, i.e. events with a tag signal in any part of the opposite side forward spectrometer (small angle tagger and large angle tagger) have been rejected, thus keeping $\Theta_{2\max} < 23 \text{ mrad}$.

(c) Three or more tracks in the central detector or 2 tracks in the central detector and at least one shower which is not associated with the tracks ($E_n > 150 \text{ MeV}$, $|\cos \Theta_n| < 0.966$). The two tracks have been defined by the following conditions: $|\cos \Theta_{h1}| < 0.629$, $p_T > 350 \text{ MeV}$, $|\cos \Theta_{h2}| < 0.866$, $p_T > 100 \text{ MeV}$, where p_T is the transverse momentum, Θ_n and Θ_{hi} are the polar angles of showers and hadrons relative to the beam axis. Two of the tracks in multitrack events are subject to the same conditions.

After applying these cuts the contamination from beam–gas scattering was approximately 15%. Using the

vertex position along the beam this background has been subtracted on a statistical basis. The resulting number of events after all cuts is 488 ± 27 .

The total cross sections σ_t and σ_q depend on Q^2 , the mass squared of the virtual photon, and on W , the invariant mass of the hadronic system. Q^2 has been determined from the energy and angle of the scattered lepton as measured in the SAT. Due to the fine granularity and the good energy resolution of the system the error in the determination of Q^2 is rather small, typically 15%. The invariant mass W of the hadronic system has been determined from the showers and tracks in the central detector assuming pion masses for all charged particles. Because of particle losses (due to the limited acceptance of our detector, reconstruction losses etc.) the visible invariant mass W_{vis} is always lower than the true mass, typically by 15–20%. Our conclusions do not depend critically on the difference between W and W_{vis} and thus an unfolding procedure has only been applied in the final step of the analysis (see below).

We have calculated the detection efficiency via a Monte Carlo program simulating the electron–photon scattering process in the PLUTO detector. The angle and energy spectra of the outgoing leptons have been generated according to the formalism discussed above [formulae (2) and (5)]. For the hadronic part of the cross section ($\sigma_t + \epsilon\sigma_q$) we have used a multipion phase space model with limited transverse momentum (as familiar from other high energy hadronic scattering processes) and constant production rate, independent of Q^2 and W . Note, that the polarization parameter is practically constant ($\epsilon \approx 0.95$) in the kinematical range discussed in this paper, and it is thus impossible to separate σ_t from σ_q .

The detection efficiency has been found to depend critically on the transverse momentum cutoff. The inclusive p_T distribution for all our events is shown in fig. 1. A single exponential does not fit the data. They are well reproduced however, if 75% of the events are generated according to $\exp(-5p_T^2)$ and 25% according to $\exp(-1p_T^2)$ (solid line in fig. 1). The charged multiplicity has been taken from low energy e^+e^- annihilation data [8–10] (parametrized by $n = 2 + 0.7 \ln W^2$) and the ratio of charged pions to neutral pions has been taken as 2 : 1. The observed charged multiplicity and the observed number of showers is well consistent with the predictions from the Monte Carlo simulation.

The resulting cross section is plotted versus W_{vis} in




Fig. 1. Inclusive p_T^2 distribution of the charged hadrons. The solid line is the prediction of the Monte Carlo simulation including the detector efficiency.




Fig. 2. The total cross section $\sigma_t + \epsilon\sigma_q$ at $\langle Q^2 \rangle = 0.25 \text{ GeV}^2$ versus the visible invariant mass. The prediction of the VMD model is given by the solid line.

fig. 2, the visible invariant mass. Fig. 3a contains $\sigma_t + \epsilon\sigma_q$ as function of Q^2 for $W_{vis} < 3.5$ GeV whereas in fig. 3b we show the same quantity for $W_{vis} > 3.5$ GeV. Fig. 3a also contains the result from ref. [5] ($\langle Q^2 \rangle = 0.1$ GeV²) measured at much smaller beam energies ($\langle E \rangle = 7$ GeV). The errors given are statistical only, we estimate an additional overall normalization error of 20% mainly due to uncertainties in the hadronic cross section model. It must be noted that in contrast to the annihilation channel the probability for the detection of two photon initiated hadronic events is only in the order of 25% (not accounting for the tagging efficiency). This detection efficiency is approximately constant above $W_{vis} = 2$ GeV and drops by a factor of two at 1 GeV. The W dependence of the data is relatively smooth, the cross section being practically constant above $W_{vis} = 3$ GeV and increasing by a factor of 2–3 towards the smallest

value of W_{vis} . It is quite obvious that the cross section drops with growing Q^2 .

For the following discussion we put $\sigma_q = 0$ in accordance with the experience from electroproduction on hadronic targets. A rather flat W dependence can be calculated from theories applying the usual tools of hadronic physics, specifically resonance–Regge duality and factorization. A “standard prediction” for $\sigma_t(0, W)$ is [11]:

$$\sigma_{\gamma\gamma}(W) = \sigma_t(0, W) = 240 \text{ nb} + 270 \text{ nb GeV}/W. \quad (6)$$

As a first guess one would assume the Q^2 dependence of the cross section to be given by the ρ -pole form factor

$$\sigma_t(Q^2, W) = \sigma_{\gamma\gamma} \cdot F_\rho^2 = \sigma_{\gamma\gamma} (1 + Q^2/m_\rho^2)^{-2}. \quad (7)$$

Both the W dependence [formula (6)] and the Q^2 dependence of $\sigma_t(Q^2, W)$ [formula (7)] can also be derived from Vector Meson Dominance, VMD.

The Q^2 dependence for $W_{vis} < 3.5$ GeV and for $W_{vis} > 3.5$ GeV is well consistent with the ρ pole as is demonstrated by the solid lines in figs. 3a and 3b which each have been normalized to the number of events in the plot. The statistics of our data is too poor to distinguish between the ρ pole and a more complicated series of vector mesons as suggested by general VMD.

We have further compared the expectation from VMD with the W dependence of our results. The solid line in fig. 2 is the expectation after folding in the detector resolution. Although at higher W_{vis} the agreement is good [$\sigma_t = (1.21 \pm 0.13) \sigma_{VMD}$ [formula (7)] for $W_{vis} > 3.5$ GeV], the VMD curve can obviously not account for the increase at smaller W_{vis} . This we have already pointed out in our previous publication [5].

We have extrapolated the data to $Q^2 = 0$ using the ρ -pole ansatz and corrected for the difference between W and W_{vis} . In doing that we have used the following ansatz for the hadronic cross section

$$\sigma_t(Q^2, W) = [A(240 \text{ nb} + 270 \text{ nb GeV}/W) + B/W^2] F_\rho^2. \quad (8)$$

The best fit to the measured invariant mass distribution was achieved for $A = 0.97 \pm 0.16$ and $B = 2250 \pm 500$ nb GeV² with $\chi^2/df = 0.5$. This result is graphically represented by the hatched band in fig. 4 where we have plotted $\sigma_{\gamma\gamma}$ versus the true invariant mass W . The limits represent the one standard deviation error of the fit.




Fig. 3. The total cross section $\sigma_t + \epsilon\sigma_q$ as function of Q^2 for $W_{vis} < 3.5$ GeV (a) and for $W_{vis} > 3.5$ GeV (b). The solid line is the prediction from ρ meson dominance.




Fig. 4. The total photon-photon cross section extrapolated to $Q^2 = 0$, versus the true invariant mass W . The hatched band represents the 1 standard deviation limits as obtained from the fit. The VMD prediction is given by the dashed line.

It has been suggested [12] to include into the theoretical predictions non-Regge terms like the quark box diagram, the magnitude of which can be very roughly estimated by (q_i is the quark charge and m_q its mass):

$$\sigma_{\gamma\gamma}^{\text{quark}} = (4\pi\alpha^2/W^2) 3 \sum_i q_i^4 \ln(W^2/m_q^2). \quad (9)$$

The most important feature of this prediction is the $1/W^2$ dependence. Including u d s quarks with masses of 100 MeV the magnitude of $\sigma_{\gamma\gamma}^{\text{quark}}$ is 650 nb at $W = 1$ GeV. Note, that for small values of Q^2/W^2 the Q^2 dependence of the bare box diagram is given by $1 - 2 \times Q^2/W^2$ which is weaker than the ρ form factor. It is thus not certain if the deviations from VMD at small W can be explained quantitatively by a simple inclusion of the box diagram [13].

In conclusion, we have measured for the first time hadron production in electron-photon scattering for

Q^2 values up to 0.6 GeV^2 and over a wide range in W . The Q^2 dependence is properly described by VMD in its simplest form (ρ pole). Above $W_{\text{vis}} = 3.5 \text{ GeV}$ the data are well consistent with the VMD prediction whereas at smaller invariant masses a $1/W^2$ term is needed which may qualitatively be understood as the contribution of point-like quarks.

We wish to thank Professors H. Schopper, G. Voss, E. Lohrmann and Dr. G. Soehngen for their valuable support. We are indebted to the PETRA machine group and the DESY computer center for their excellent performance during the experiment. We gratefully acknowledge the efforts of all engineers and technicians of the collaborating institutions who have participated in the construction and maintenance of the apparatus.

References

- [1] Ch. Berger, in: Lecture notes in physics, Vol. 134 (Springer); PLUTO Collab., Ch. Berger et al., DESY 80/94.
- [2] V.M. Budnev, I.F. Ginzburg, G.V. Meledin and V.G. Serbo, Phys. Rep. 15C (1975) 182.
- [3] C. Carimalo, P. Kessler and J. Parisi, Phys. Rev. D21 (1980) 669.
- [4] S.J. Brodsky, T. Kinoshita and H. Terazawa, Phys. Rev. D4 (1971) 1532.
- [5] PLUTO Collab., Ch. Berger et al., Phys. Lett. 89B (1979) 120.
- [6] J.H. Field, Nucl. Phys. B168 (1980) 477.
- [7] PLUTO Collab., Ch. Berger et al., Phys. Lett. 81B (1979) 410.
- [8] C. Bacci et al., Phys. Lett. 86B (1979) 234.
- [9] J.L. Siegrist, Ph. D. thesis, SLAC-PUB-225 (1980); UC-34d (1980).
- [10] A. Bäcker, Ph. D. thesis, Universität Siegen, Internal report DESY F33 77/03 (1977).
- [11] T.F. Walsh, J. de Phys. C2 Suppl. 3 (1974) 77; S.J. Brodsky, J. de Phys. C2 Suppl.3 (1974) 69; J.L. Rosner, in: ISABELLE Physics Prospects, BNL report 17552 (1972) 316.
- [12] M. Greco and Y. Srivastava, Nuovo Cimento 43A (1978) 88; S.J. Brodsky, F.E. Close and J.F. Gunion, Phys. Rev. D6 (1972) 177; T.F. Walsh and P. Zerwas, Phys. Lett. 44B (1973) 195.
- [13] J. Gunion, Invited talk Intern. Workshop on Photon-photon collisions (Amiens, 1980), SLAC-PUB-2503 (1980).